

# Screening Some Accessions Of Lentil (*Lens Culinaris* M.) For Salt Tolerance At Germination And Early Seedling Stage In Eastern Ethiopia

Azene Tesfaye, Yohannes Petros, Habtamu Zeleke

Arba Minch University, Biodiversity Research Centre,  
Haramaya University, Department of Biology, P.O.Box.138, Dire Dawa, Ethiopia  
Haramaya University, School of Plant Science, P.O.Box.138, Dire Dawa, Ethiopia.  
E-mail: azet567@gmail.com /+251921574929

**Abstract:** To evaluate genetic variation among Ethiopian lentil, laboratory experiment were conducted to screen 12 accessions of lentil (*Lens culinaris* M.) for salt tolerance. Seeds of 12 Lentil accessions were grown at laboratory (Petri dish) condition with different levels of salinity (0, 2, 4, and 8 dSm<sup>-1</sup> NaCl) for 4 weeks. The experimental design was completely randomized design (CRD) in factorial combination with three replications. Data analysis was carried out using SAS software. Average germination time, germination percentage, seedling shoot and root traits, seedling shoot and root weight were evaluated. The two way ANOVA for varieties revealed statistically significant variation among lentil accession, NaCl level and their interactions ( $p < 0.001$ ) with respect to the entire parameters. It was found that salt stress significantly delays germination rate and decreases germination percentage, shoot and root length, seedling shoot and root weight of lentil accessions. The degree of decrement varied with accessions and salinity levels. Accessions 36120, 9235 and 36004 were better salt tolerant than the other accessions. As the result, it is recommended to be used as a genetic resource for the development of lentil accession and other very salt sensitive crop with improved germination under salt stress condition.

**Key words:** Accession, Germination, Lentil, NaCl, Screening, Seedling stage

## Introduction

Salinity is one of the most serious factors that hamper the productivity of agricultural crops, with adverse effects on germination, plant vigor and crop yield [33] particularly in arid and semi-arid regions of the world [3] because of the lack of sufficient amount of rainfall leads to leaching the accumulated salt and also it affects many irrigated areas mainly due to the use of underground water. Salt-affected soils are distributed throughout the world and no continent is free from the problem [14]. In Ethiopia, salt-affected soils are prevalent in the Rift Valley and the lowlands [26]. Salinity stress cause a multitude of physiological problems in plant processes [32]. It causes a significant reduction in germination percentage, germination rate, shoot and root length, root and shoot weight, and dry root and shoot weight, and seed yield which lead to the death of the entire plant [20]- [23]. Germination and seedling growth under saline environment are the screening criteria that are widely used to select the salt tolerance genotype [7] [10]. Because of salinity tolerant at this stage was shown to be a heritable trait that enable the crop salt tolerant throughout its growth stage [10] and [11] although it is a polygenic character linked to a complex genetic basis [28] and seeds and young seedlings are frequently was affected by much higher salinities than vigorously growing plants because germination usually occurs in surface soils, which accumulate soluble salts because of evaporation and capillary rise of water [5]. Since grain legumes especially lentil are salt sensitive, farmers do not consider growing them in a saline environment, though; there is a considerable difference in salt tolerance among crops/accessions [3]. Screening of available Lentil accessions is important to find a relative salt tolerant accession. Therefore, the general objective of this study was to assess the genetic variability for salinity tolerance among some lentil accessions, specifically to evaluate the effect of salt on germination and early

seedling stage of Lentil accessions and to identify salt tolerant Lentil accessions.

## Methods and material

### Description of the study area and plant material

The experiment was conducted in the Botanical science laboratory, Department of Biology, Haramaya University, Ethiopia. Seeds of twelve Lentil accessions were obtained from the Ethiopia Institute of Biodiversity (EIB).

**Table1.** Descriptions of lentil accessions that were used in the experiment

Accession Code	Region/ State/	Zone	Woreda/ District	Latitude	Longitude	Altitude
9235	Oromiya	MisrakHarerge	Meta	09-16-21-N	41-33-45-E	2535
36004	Amara	Semen Shewa	Ankober	09-39-00-N	39-41-00-E	3180
36006	Oromiya	MisrakShewa	Gimbichu	08-57-00-N	39-05-00-E	2370
36019	Oromiya	MirabShewa	AlemGena	08-48-00-N	38-20-00-E	2150
36025	Gumuz	Metekel	Wenbera	Unknown	Unknown	1580
36032	Oromiya	Bale	Ginir	Unknown	Unknown	1520
36064	SNNP	Bench Maji	Dirashe	Unknown	Unknown	Unknown
36093	Oromiya	MirabHarerge	Chiro	09-04-00-N	40-41-00-E	2000
36094	Oromiya	MirabHarerge	Chiro	09-02-00-N	40-44-00-E	1870
36095	Somali	Shinile	Afdem	Unknown	Unknown	1800
36113	Oromiya	MisrakHarerge	Deder	Unknown	Unknown	Unknown
36120	Oromiya	MirabWellega	Gawo Dale	Unknown	Unknown	1870

**Treatments and Experimental Design:** The study was conducted under laboratory condition at room temperature and based on morphological variation among lentil accessions in order to assess the salt tolerance in terms of seed germination and seedling growth. Four different NaCl solutions with salinity levels of 0, 2, 4, and 8 dS/m [8] and [31] were prepared by dissolving 0, 1.28, 2.56 and 5.12 gm. of NaCl in one liter of water respectively. 12 accession of Lentil. The experiment was laid as a Completely Randomized Design (CRD) in a factorial arrangement and replicated three times. The treatments were assigned randomly to each Petri dish.

**Experiment procedure:** In order to assess the response of the 12 Lentil accessions under different concentration of NaCl, 12 seeds were first surface sterilized in 5 % sodium hypochlorite solution for 20 minute and washed three times with sterilized distilled water. Prior to experiment 10 cm diameter Petri dishes were thoroughly washed and sterilized in hot air oven at 70 °C for 36 hours and Whatman filter paper for 24 hours at 70 °C [26]. After sterilization, Petri dishes were lined with Whatman No.3 filter paper and treated with 10 ml of deionized water (control), 2, 4, and 8 dS/m of NaCl. Following this, twelve uniform seeds of each Lentil accessions were placed on each Petri dish approximately in uniform distance. The Petri dishes were arranged in a completely randomized design (CRD) in a factorial combination with three replications [19]. Each Petri dish was treated with 10 ml of the respective concentrations of NaCl in every other day. Salt levels were maintained each day by dripping out and applying fresh salt solution. The Petri dishes were put within a glass box to avoid loss of moisture through evaporation. Germination started after two days of sowing and the germination count was continued until the 9<sup>th</sup> day. Germination was recorded daily and a seed was considered germinated when both plumule and radicle had emerged  $\geq 0.05$  cm [2].

**Germination rate:** the average number of days needed for plumule or radical emergence calculated following the formula described by [26].

**Germination percentage:** Seven days after seeds were put into the Petri dishes, six germinated seeds were

counted, and the germination percentage calculated. The germination percentage calculated [6] as:

$$\text{Germination percentage} = \frac{\text{Number of seeds germination}}{\text{Number of seeds sown}} \times 100$$

**Salt tolerance index** was calculated as total plant (shoot + root) dry weight obtained from 6 randomly selected seeds grown on different salt concentrations compared to total plant dry weight obtained on normal concentration. {[STI = (TDW at Sx/TDW at S1) x 100], STI= salt tolerance index, TDW = total dry weight, S1 = control treatment, Sx = x treatment}[25].

**Seedling Shoot Length (cm):** Fifteen days after germination, shoot length of 6 randomly picked seedlings from each Petri dish measured in centimeters [18].

**Seedling Root Length (cm):** Fifteen days after germination, root lengths of 6 randomly picked seedlings from each Petri dish were measured in centimeters.

**Seedling Shoot-to-Root Ratio:** calculated as the ratio of seedling shoot length to seedling root length [25].

**Seedling Fresh Shoot Weight (g)** was measured after 30 days of sowing by weighting the mass of shoots of 6 randomly picked seedlings from each Petri dish using sensitive balance.

**Seedling Fresh Root Weight (g):** measured by weighting the mass of roots of six randomly picked seedlings after 30 days of sowing from each Petri dish using sensitive balance.

**Seedling Shoot Dry Weight (g):** measured by picking six seedlings randomly from each petri dish and oven drying their shoots at 80 °C for 48 hours and weighting them using sensitive balance.

**Seedling Root Dry Weight (g):** measured by picking six seedlings randomly from each Petri dish after 30 days of sowing and oven drying their roots at 80 °C for 48 hours and weighting them using sensitive balance [25].

#### Data analysis

The data were subjected to analysis of variance using SAS (version 9.1) and the means were separate using the

Least Significant Difference (LSD) test at 5% significance level.

## RESULTS AND DISCUSSIONS

### Influence of Salinity on Germination of Lentil Accessions

**Germination rate** Two-way analysis of ANOVA for germination rate showed that the NaCl salt levels highly significantly ( $p < 0.001$ ) influenced germination rate. Nevertheless, accessions responded differently to different salinity levels. For instance, at  $2 \text{ dSm}^{-1}$  salinity level, accession 36120, 36004 and 9235 had the shortest germination rate. Thus, those accession had needed shorter time to be germinated than the other accessions while, accession 36019, 36032 and 36064 attained the longest germination rate and needed longer time to be germinated (fig 1). Moreover, at 4 and  $8 \text{ dSm}^{-1}$  salinity level, accession 36120, 9235 and 36004 had the shortest germination rate and germinated faster than other accessions that were tested (fig 1). Whereas, accession 36019, 36064 and 36093 had longer germination rate and needed more time to be germinated (fig 1). Accessions 36120, 9235 and 36004 germinated faster than the other accession at 2, 4 and  $8 \text{ dSm}^{-1}$  salinity level whereas, salinity delayed seed germination of accession 36019, 36093, and 36025 and those accessions needed longer time to germinated in the entire salt treatments (fig 1). The result revealed that salinity delays the germination rate of lentil accessions and the result was in full agreement with the previous studies of [3] and [15] in lentil; [24] in *Phaseolus* species who reported that salinity delay seed germination and decrease germination rate. This might be due to salinity affects germination by facilitating intake of toxic ions and hinder many metabolic, physiological, and enzymatic activities. In addition, salinity may cause osmotic potential which result decreasing absorption of water.




fig 1. Effect of salinity on germination rate of lentil accessions

**Germination percentage** the two-way analysis of variance (ANOVA) found highly significant variation in germination percentage among accessions, salinity and their interaction ( $p < 0.001$ ). At  $2 \text{ dSm}^{-1}$  salinity (NaCl) level, accession 9235, 36004 and 36120 performed well and achieved the higher percent of germination whereas, salinity highly

affected seed germination of accession 36019 (fig 2). Moreover, at 4 and  $8 \text{ dSm}^{-1}$  salinity level, the highest value of germination percentage was recorded in accession 9235, 36120 and 36004 whereas, the lowest value was recorded in accession 36019 and 36094 (fig 2). The result indicates that the percent of germination generally decreased with increasing salt concentration. The findings of the study were in line with previous research studies reported by [35] in *P.sativum* and [3] in lentil who reported that a significant decrease in seed germination were observed while increasing salinity level. The reason is assumed to be due to salinity that attributed to osmotic retention of water (water deficiency).




fig 2. Effect of salinity on seed germination of lentil accessions

**Salt tolerance index** Analysis of variance (ANOVA) for salt tolerance index showed that salt tolerance index was highly significant difference among accession, salinity level, and accession\*salinity interaction ( $p < 0.001$ ). Even though at  $2 \text{ dSm}^{-1}$  salinity level, salinity reduced the overall growth of accession 9235, 36004 and 36120, the degree of reduction was lesser on those accession, hence, the maximum salt tolerance index were observed at those accession while, salinity highly reduced the overall growth of accession 36019. Thus, accession 36019 exhibited the minimum percent of salt tolerance index than the rest of the accessions (fig 3). Moreover, at 4 and  $8 \text{ dSm}^{-1}$  salinity level, salinity easily hampered the overall growth of accession 9235, 36120 and 36004 than the other accession while, the overall growth of accession 36019 were highly hindered by salinity and as result; this accession attained the minimum value of salt tolerance index (fig 3). The result indicates that salt tolerance index of lentil accession were significantly reduced as salinity concentration increased (fig 3). The result was in line with previous studies of [25] who reported the salt tolerance index of lentil decrease when salinity levels become increased.




fig 3. Effect of salinity on salt tolerance of lentil accessions

**Influence of Salinity on Seedling Shoot and Root traits of Lentil Accessions**

**Seedling Shoot Length (SSL) (cm)** The analysis of variance (ANOVA) for seedling shoot length data showed that highly significant variation in seedling shoot length ( $p < 0.001$ ) among accessions, salinity level and their interaction. Seedling shoot length of lentil accessions varied among accessions and salinity level (fig 4). At 2  $dSm^{-1}$  salinity level, accession 36120 followed by accession 9235 and 36004 attained the longest shoot length while, accession 36094 had the shortest shoot length (fig 4). Moreover, accession 36120, 9235 and 36004 performed well and attained the longest shoot length at 4 and 8  $dSm^{-1}$  salinity levels (fig 4) whereas, the shortest shoot growth was observed in accession 36019, 36094 and 36113 (fig 4). This result showed that increment of NaCl treatments resulted in a significant reduction in shoot growth (fig 4). The findings of this studies show conformity with research result of [25] and [27] who reported that salinity inhibits elongation of shoot in lentil and increasing NaCl treatment results significant reduction of shoot growth. The reduction in shoot length probably because genetic variation among lentil accession and excessive accumulation of salts in the cell wall elasticity, thus, secondary cell appears sooner and cell wall becomes rigid as a consequence the turgid pressure efficiency in cell enlargement decreases that result in short shoot growth




fig 4. Effect of salinity on seedling shoot length of lentil accessions

**Seedling Root Length (SRL)** The analysis of variance (ANOVA) for seedling root length indicates that there was highly significant variation in seedling root length among lentil accessions, salinity levels and their interaction ( $p$

$< 0.001$ ). At 2  $dSm^{-1}$  salt concentration, accession 36120, 9235 and 36004 attained the longest root length than other accession while, salinity adversely reduced root growth in accession 36094 and this accession had shortest root length (fig 5). Furthermore, at 4 and 8  $dSm^{-1}$  salt concentration, the longest root length was recorded in accession 36120, 9235 and 36004 than the rest of the accessions whereas; the shortest root length was observed in accession 36094, 36113, 36093, and 36019 (fig 5). The result elucidate that the increment of NaCl concentration was cause the reduction of seedling root length of lentil accession (fig 5) The result is in full agreement with [10] in lentil; [7] in senna plant; [16] in pea who reported that high salinity reduced root length in lentil.




fig 5. Effect of salinity on seedling root length of lentil accessions

**Shoot to Root length Ratio (SRR)** The two-way analysis of variance (ANOVA) for seedling shoot-to-root length ratio (SRR) showed significant variations for accessions, salinity level and accession \* salinity interaction ( $p < 0.001$ ). At 2  $dSm^{-1}$  salinity level, accession 36093 and 36095 had the lower value of shoot to root ratio than the other accessions (fig 6). Moreover, at 4  $dSm^{-1}$  NaCl level, accession 36064 followed by 36025 and 36095 showed the higher seedling shoot to root ratio. However, accession 36019 attained significantly lower values seedling shoot to root ratio than the other accession (fig 6). At 8  $dSm^{-1}$  NaCl level, accession 36120, 36004 and 9235 attained significantly higher mean seedling shoot to root ratio in contrast to this, accession 36019 showed the lowest value (fig 6). The result indicates that some accessions show significantly higher reduction in SRR as increment of NaCl concentration in the growth media. In contrast to this decrement of SRR in some accession were observed as the salinity level increased. The findings of this study showed conformity with the previous studies on other legumes [26] in haricot bean, [1] in faba bean who reported that SRR was highly reduced at higher salinity levels for some accessions and the accessions showed significant variation in their response to salinity.




fig 6. Effect of salinity on seedling shoot to ratio of lentil accessions




fig 7. Effect of salinity on seedling fresh shoot weight of lentil accessions

**Seedling fresh Shoot weight (SFSW) (g)** Two way analysis of ANOVA seedling shoot fresh weight confirmed that there were highly significant differences among all accessions, NaCl treatments and their interaction ( $p < 0.001$ ). The NaCl treatments used in caused significant reduced in seedling fresh shoot weight. Some accession attained the maximum value of seedling fresh shoot weight while other attained the minimum value of seedling fresh shoot weight. For instance, accession 9235, 36120 and 36004 attained the longest shoot length and produced the maximum value of seedling fresh shoot weight at 2, 4 and 8  $dSm^{-1}$  salinity level (fig 7) while, minimum value of seedling fresh shoot weight was recorded in accession 36019, 36094, and 36093 (fig7). Salinity reduced shoot growth of all accession as compared to the control but the degree of reduction was varied between accessions and salt concentration. For instance, accession 36120, 9235 and 36004 attained relatively the maximum value of seedling fresh shoot weight even at higher salinity level than the other accessions and considered as salt tolerant while, salinity adversely reduced shoot growth of accession 36019, 36094, and 36093. As result, those accessions exhibited the minimum value of seedling fresh shoot weight even at lower salt concentration (fig 7). The result showed that the seedling fresh shoot weights of lentil accessions significantly reduced with increment salinity level (fig7). The result was in line with previous research findings of [38] who reported as that there was a rapid decrease in seedling fresh shoot weight of leguminous plants under saline environment. This reduction may be due to limited supply of metabolites to young growing tissues, because metabolic production takes place within in the leaves and is significantly perturbed at high salt stress, either due to the low water uptake or toxic effect of NaCl concentration [21],[32]-[40].

**Seedling root fresh weight (SRFW) (g)** Two way analysis of ANOVA for seedling fresh root weight confirmed that there were highly significant differences among lentil accessions NaCl treatments and accessions \* treatment interaction ( $p < 0.001$ ). At 2  $dSm^{-1}$  salinity level, accession 9235 followed by 36120 and 36004 achieved the maximum value of seedling fresh root weight whereas, the minimum value of seedling fresh root weight was recorded in accession 36019 (fig 8). Moreover, at 4 and 8  $dSm^{-1}$  some accession performed well and attained the maximum value of seedling fresh root length while, other accession had the minimum value of seedling fresh root weight. For instance, accession 36120, 9235 and 36004 achieved the maximum value of seedling fresh root weight than the other accession whereas; accession 36019 attained the minimum value of seedling fresh root weight (fig 8). Salinity reduced growth of root in accession 36120, 36004 and 9235 but; the degree of reduction of seedling fresh root weight in those accessions was less than the other accessions. Consequently, those accessions had the maximum value of seedling fresh root weight even at higher salinity level than the other accessions. On the other hand, salinity highly inhibited the root elongation of accession 36019 and this accession had minimum value of the seedling fresh root weight. The result justified that salinity reduced seedling fresh root weight of lentil accessions (fig 8). The findings of this result in line with the result of findings of [25] and [24] who reported salinity increment significantly reduced fresh root weight in Lentil and phaseolus species, respectively.




fig 8. Effect of salinity on seedling fresh root weight of lentil accessions

**Seedling Dry Shoot Weight (SDSW) (g)** Statistical analysis for seedling shoot dry weight revealed that there were highly significant differences among all accessions, NaCl treatments and accessions \* treatment interaction ( $p < 0.001$ ). At  $2 \text{ dSm}^{-1}$  salinity level, accession 9235, 36120 and 36004 achieved maximum value of seedling dry shoot weight than the other accession (fig 9) whereas, the minimum value of seedling dry shoot weight was scored in accession 36019 (fig 9). Moreover, at 4 and  $8 \text{ dSm}^{-1}$  salt concentration, accession 9235 and 36120 achieved the maximum value of seedling dry shoot weight than the rest of the accessions that were tested while, accession 36019 and 36113 attained the minimum value of seedling dry shoot weight (fig 9). This result revealed that salinity antagonistically reduced in seedling dry shoot weight (fig 9a&b). The result show full agreement with previous research result reported of [41] and [22] on lentil; [39] in rice and [13] in phaseolus species who reported that increasing the concentration of salinity cause significant reduction in shoot growth consequently reduced seedling dry shoot weight.




fig 9. Effect of salinity on seedling dry weight of lentil accessions

**Seedling Root Dry Weight (SRDW) (g)** Analysis of variance for seedling dry root weight exhibited that highly significant variations among NaCl treatments, accessions and their interaction ( $p < 0.001$ ). accession 9235, 36120 and 36004 had the maximum weight of seedling dry root weight than the other accession that were tested at 2, 4 and  $8 \text{ dSm}^{-1}$  salinity level, (fig 10), in contrast to this, accession 36095, 36019 and 36025 attained the minimum value of seedling dry root weight (fig 10). The result showed that significantly reduction in mean root dry weight when the salinity concentration increase (fig 10). This result showed analogous to earlier studies [7], [9] and [12] on Lentil [4] on *Gossypium hirsutum*, who reported that salt stress caused a significant decrease dry weight of root tissues.




fig 10. Effect of salinity on seedling dry root weight of Lentil accessions

### conclusion

Salinity is a continuing problem in the arid and semi-arid tracts of the world. It could be alleviate during irrigation management and/or crop management. However, the former approach is outdated and very expensive. Nevertheless, the latter is economical as well as efficient and it enables to produce salt tolerant crop lines. However, prior to that there is a need to confirm the presence of genetically based variation for salt tolerance among different species or varieties of a particular crop at different growth stages. The presence of genetic variation offers a basic tool for evaluating effect of salinity on lentil accessions and to overcome the presence of large number of variation for relatively salt tolerant lentil accession and it will appreciated to find accession with gene tolerant to salinity. Screening of salinity tolerance under field condition involves many environmental factors that affect genetic and phenotypic expression of accessions. Hence, controlled environment, Laboratory and greenhouse screening method indicate to be an ideal method to screen large amount of accessions with less efforts and accurately. Thus, the correct and clear expression of Lentil accessions for salt tolerant can be evaluated by this method using different NaCl level. The findings of this work confirmed that response of lentil accession to salinity show significant variation as their expose to different salinity level. The result explain that most out that all of the morpho-physiological and yield and yield related traits considered were significantly decreased with higher levels of salinity. Out of twelve lentil accession, accession 36120, 36004 and 9235 performed well under salt stress conditions in most of the parameter for both laboratory and greenhouse experiment as result those accession were recommended to be sown in saline condition.

### Acknowledgment

Right from the outset, I would like to thank the supreme Almighty God for providing me the patience and endurance to complete the study. The author is especially indebted to the thesis research advisors Dr. Yohannes Petros, Dr. HabtamuZelege and Mr. chalew Tesfaye for their indispensable support, scholarly guidance and critical suggestions, inspiration, encouragement throughout the study period, which were highly instrumental in enabling him not only to complete his study smoothly and successfully but also to make the undertaking profoundly

educational. My special thanks go to Haramaya University for providing institutional support.

## References

- [1] Abdelhamid, M. T., M. B. Shokr and M.A. Bekheta, 2010. Effects of induced salinity on four vicia faba cultivars differing in their broomrape tolerance. pp 12-18. Fourteenth International Water Technology Conference, IWTC 14, Cairo, Egypt.
- [2] Abdul W.I., Ahmad H., Ghulam Q., Ghulam M., M Tariq and A Muhammad. 2006. Effect of Salinity on Germination, Growth, Yield, Ionic Balance and Solute Composition of Pigeon Pea (*Cajanus cajan* (L.) Millsp. Pak. J. Bot., 38(4): 1103-1117.
- [3] Ahmed, S., 2009. Effect of soil salinity on the yield and yield components of Mung bean. Pak. J. Bot., 4(1): 263-268.
- [4] Akhtar, J. and F.M. Azhar, 2001. Response of *Gossypium hirsutum* L. Hybrids to NaCl salinity at Seedling Stage. IJAB 03 (2): 233-235.
- [5] Almansouri, M., J.M. Kinet and S. Lutts, 1999. Compared effects of sudden and progressive impositions of salt stress in three durum wheat (*Triticum durum* Desf.) cultivars. J. Plant Physiol., 154: 743-752.
- [6] AniatulHaq.R. Vamil and R.K., Agnihotri, 2012. Effect of Osmotic Stress (PEG) on Germination and Seedling Survival of Lentil (*Lens culinaris* M. Res. Agri. Sci. 1(3): 201-202.
- [7] Arshi, A., A.Z. Abdin, and M. Iqbal, 2003. Growth and metabolism of senna as affected by salt stress. Biol. Plant. 45:295-298.
- [8] Asghar, R., N. Rob, Mc. David and M.H. Shahab, 2009. Effects of Salinity and Temperature on Germination, Seedling Growth and Ion Relations of two Lentil (*Lens culinaris*) Cultivars. J. Seed Techno, 31(1):76-86.
- [9] Ashraf, M & McNeilly, T., 1988. Variability in salt tolerance of nine spring wheat cultivars. J. Agro. Crop Sci, 160:14-21.
- [10] Ashraf, M. and Waheed, A. 1990. Screening of local/exotic accessions of lentil (*Lens culinaris*) for salt tolerance at two growth stages. Plant Soil 128:167-176.
- [11] Ashraf, M., 1994. Organic substances responsible for salt tolerance in *Eruca sativa*. J. Biol. Plant., 36:255-259.
- [12] Badeoglu, E., F. Eyidogan, M. Yucel and H.A. Oktem, 2004. Antioxidant responses of shoots and roots of lentil to NaCl. Pak. J. Bot., 43: 269-274.
- [13] Bayuelo Jimenez, J.S., D.G. Debouck and J.P. Lynch, 2002. Salinity tolerance in *Phaseolus* species during early vegetative growth. Crop Sci., 42: 2148-2192.
- [14] Brady, N.C. and R.R. Weil, 2002. The Nature and Properties of Soils, 13<sup>th</sup> Edition, Prentice- Hall, Upper Saddle Rivers, New Jersey. 11 p.
- [15] Carter, C.T., C.M. Grieve and J.P. Poss, 2005. Salinity effects on emergence, survival, and ion accumulation of *Limonium perezii*. J. Plant Nutr, 28:1243-1257.
- [16] Duzdemiro., A. Kurunc And A. Unlukara, 2009. Response of Pea (*Pisum sativum*) To Salinity and Irrigation Water Regime. Bulgarian Journal Of Agricultural Science, 15 ( 5) 2009, 400-409.
- [17] EL-Khashab, A., A.M. Elaidy, A.A. EL-Sammak, .A. F. Salama, and M.I. Rienger, 1997. Paclobutrazol reduces some negative effects of salt stress in peach. J. Amer. Soc. Hort. Sci., 122(1): 43-46.
- [18] Geressu Kinfe Michael and Gezaghegne Melkamu, 2008. Response of some lowland growing sorghum (*Sorghum bicolor* Moench L.) accessions to salt stress during germination and seedling growth. Afr. J. Agric. Res., 3 (1): 44-48.
- [19] Gomez, K.T. and A.A. Gomez, 1984. Statistical Procedures for Agricultural Research, 2<sup>nd</sup> Edition. John Wiley and Sons, New York.
- [20] Gupta R., and P S. Minhas, 1993. Managing salt affected waters for crop production. Pp: 159-198. In: Singh S.D. (Ed.), Arid Land Irrigation and Ecological Management. Scientific Publishers, New Delhi
- [21] Hussain, K., A. Majeed, K. Nawaz, K.H. Bhatti and F.K. Nisar, 2009. Effect of different levels of salinity on growth and ion contents of black seeds (*Nigella sativa* L.). Curr. Res. J. Biol. Sci., 1(3): 135-138.
- [22] Islam M.T., N.A. Jahan, A.K. Sen and M.H.R. Pramanik, 2012. Effects of Salinity On Morpho-Physiological attributes and Yield of Lentil Genotypes Int. J. Sustain. Crop Prod. 7(1):12-18.
- [23] Jamil, M., D.B. Lee, K.Y. Jung, M. Ashraf, S.C. Lee and E.S. Rha, 2006. Effect of salt (NaCl) stress on germination and early seedling growth of four vegetables species. J. Gen. Europ. Agri., 7(2): 273-282.
- [24] Jeannette, S., R. Craig and J.P. Lynch, 2002. Salinity tolerance of *Phaseolus* species during germination and early seedling growth. Crop Sci., 42: 1584-1594.
- [25] Kagan, K., T. Karakoy, A. Bakoglu 3 and M. Akçura, 2010. Determination of salinity tolerance of some lentil (*Lens culinaris* M.) varieties. J. Food. Agri. Environ 8 (1) 140-144.
- [26] Kinfe Michael Geressu, 2011. The Response of Some Haricot Bean (*Phaseolus vulgaris*) Varieties for Salt

- Stress during Germination and Seedling Stage. *Curr.Res. J. Biol Sci.*3(4): 282-288.
- [27] Mane, A. V., G. D. Saratale, B. A. Karadge, and J. S. Samant, 2010. Microstructure, physicochemical properties and in vitro digestibility of starches from different Indian lentil (*Lens culinaris*) cultivars. *Carbohydrate Pol.*, 79: 349–350.
- [28] Mano, Y. and Takeda, K. (1997): Mapping quantitative trait loci for salt tolerance at germination and seedling stage in barley (*Hordeum vulgare* L.). *Euphytica* 94, 263-272.
- [29] Marler, T. E and M. V Mickelbart, 1993. Growth and chlorophyll fluorescence of *Spondias Purpurea* L. as influenced by salinity. *J. Trop. Agri.*, 70(3): 245-247.
- [30] Mehmood, E., R. Kausar, M. Akram and S.M. Shahzad. 2009. Is boron required to improve rice growth and yield in saline environment. *Pak. J. Bot.*, 41(3):1339-1350.
- [31] Mohammad k. G., 2012. Effectiveness of nutrient management in managing saline agro- ecosystems: a case study of *lens culinaris* medik. *pak. J. bot.*, 44: 269-274.
- [32] Munns, R., 2002. Comparative physiology of salt and water stress. *Plant Cell and Environment*, 25(2), 239-250.
- [33] Munns, R., and M. Tester, 2008. Mechanisms of salinity tolerance. *Annual Review of Plant Biology*, 59: 651-681.
- [34] Nasir Khan, M., Siddiqui, M. H., Mohammad, F., Masroor, M., Khan, A. and Naeem, M. (2007). Salinity induced changes in growth, enzyme activities, photosynthesis, proline accumulation and yield in linseed genotypes. *World J. Agric. Sci.*, 3: 685.
- [35] Noreen, Z., M. Ashraf and M. UI-Hassan, 2007. Inter-accessional Variation for Salt Tolerance in Pea (*Pisum Sativum* L.) at Germination and Screening Stage. *Pak. J. Bot.*, 39(6): 2075-2085.
- [36] Parida, A.K. and A.B. Das, 2005. Salt tolerance and salinity effects on plants. *A Rev. Ecotoxicol. Environ. Safety*, 60: 324-349.
- [37] Patil, S.L., C.S. Hunshal, B.T. Nadagouda, R.N. Kori, S.R. Salakinakop, 1996. Dry matter accumulation in lentil as influenced by saline water irrigation. *Adv. Agril. Res. India*, 6:78-87.
- [38] Stoeva, N., and M. Kaymakanova, 2008. Effect of Salt Stress on the Growth and Photosynthesis Rate of Bean Plants (*Phaseolus Vulgaris* L.). *J. Central Europ. Agri.* 9(3): 385-392.
- [39] Shereen A., R. Ansari, S. Raza, S. Mumtaz, M.A. Khan and M. Ali Khan, 2012. Salinity Induced Metabolic Changes In Rice (*Oryza Sativa* L.) Seeds during Germination. *Pak. J. Bot.* 43(3): 1659-1661, 2011.
- [40] Taffouo, V. D., J. K. Kouamou, M. T. Ngalangue, A. N. Ndjedji and A. Akoa, 2009. Effects of Salinity Stress on Growth, Ions Partitioning and Yield of Some Cowpea (*Vigna unguiculata* L. Walp.) Cultivars. *Intl. J. Bot.*, 5: 135-143.
- [41] Turan, M. A., N. Turkmen and N. Taban, 2007. Effect of NaCl on stomatal resistance and proline, chlorophyll, Na, Cl and K concentrations of Lentil Plants. *J. Agron.*, 6: 378-381.